

Organic Geochemistry Characteristics of Aliphatic Hydrocarbon Fraction of Crude Oil from Tarakan Basin, North Borneo, Indonesia

Y. Zetra, M. Fitriyyah, R.Y. P. Burhan, Z. V. Nugraheni

Department of Chemistry,
Institut Teknologi Sepuluh Nopember, Surabaya, 60111
e-mail: yzetra@gmail.com

Abstract— The study of organic geochemistry of crude oil from Pamusian field, Tarakan Basin, North Borneo has been done. The oil is fractionated by column chromatography and identified using gas chromatography-mass spectrometer (GC-MS). The presence of long chain *n*-alkanes, cadinane, 4 β (H)-eudesmane, and 18 α (H)-oleanane indicates organic matter derived from resin dammar Angiospermae family Dipterocarpaceae. It is also reported that there are drimane together with homodrimane and hopane as an indicator of bacterial input. The LHCPI value of 2.03 also indicates a high input of photosynthetic bacteria. Pr/Ph ratio of 3.76 and a drimane/homodrimane ratio of 1.058 indicating the oxic depositional environment of the sample. Isomer analysis of 18 α (H)-oleanane and 17 α (H),21 β (H)-hopane provides information that the crude oil from Tarakan Basin is mature.

Keywords—Tarakan Basin Crude Oil, Organic Geochemistry, Aliphatic Hydrocarbon Biomarker, GC-MS

I. INTRODUCTION

CRUDE oil needs by Outlook Energy Indonesia (2018) is expected to increase more than 3 times in the period 2016-2050 with an average growth of 3.3% per year [1]. This condition causes the possibility to import crude oil even higher. However Indonesia still has the potential to increase the production of crude oil and condensate, where there are still many basins that have the potential of oil reserves scattered throughout Indonesia. More than 60 sedimentary basin in Indonesia, some of which have been proven to be productive as oil and gas producers. The basin is largely located on the island of Sumatra and Borneo. One of the still-productive basins is the Tarakan basin [2].

The Tarakan Basin is located in the east part of Borneo and is divided into 4 sub-basin namely Tidung sub-basin, Tarakan sub-basin, Berau sub-basin, Muara sub-basin [3]. It is bounded by the Sekatak Berau Ridge on the west, the Suikerbrood Ridge and the Mangkalihat Peninsula to the south, the Sempurna ridge in the north, and the Sulawesi Sea in the East [3]. Oil exploration in the Tarakan Basin was conducted since 1899 and has been found 14 oil and gas fields. Up to now 86% of oil production comes from two fields, namely Pamusian field in Tarakan Island and Bunyu field on Bunyu Island [4].

Oil production of Pamusian field in Tarakan sub-basin has entered the drain stage [4]. The reactivation of abandoned oil wells is one of the attempts to increase crude oil production [5]. But this effort requires a large fee so that the feasibility data is required for well before reactivation [6]. The exploration and production of petroleum can be improved,

one of them through the application of organic geochemical sciences [7] as a complement to geological data [8]. Geochemical characteristics can be used to determine the relationship between petroleum and source rocks, mapping the geographical conditions of the petroleum system and rock source, as well as assess the timing, migration, and accumulation of oil [7].

To reactivate the well, it is necessary to know the hydrocarbon potential of the well. The maturity and source of the main organic material has a significant effect on the potential of hydrocarbons and the productivity of a source rock [9,10,11]. The source of the dominant organic material that provides good hydrocarbon potential is the source of organic material type 3 or derived from higher plants [11]. Hu et al. (2019) has used biomarkers to determine the hydrocarbon potential of Guangdong province, China. Some of the essential biomarker as indicators are long chain *n*-alkanes (*n*-C₂₇, *n*-C₂₉, *n*-C₃₁), isoprenoid C₁₉ and C₂₀, tricyclic terpane and C₂₉ Sterane.

The study of organic geochemistry through biomarker analysis has been conducted by researchers on sedimentary, such as petroleum and coal [12,13,14]. Some research through biomarker analysis of the aliphatic hydrocarbon fraction has been reported to identify the source of organic matter, the depositional environment, and the thermal maturity of petroleum [12,13,14]. Biomarker such as alkanes, isoprenoid, terpene, and sterane are groups of aliphatic hydrocarbon compounds found in petroleum [6,15,16]. The loss of *n*-alkane, isoprenoid, terpane, and sterane during the secondary process can be used to determine the degree of biodegradation in the reservoir [6] and determine the level of oil maturity [17]. The comparison of pristane to phytane and the distribution of *n*-alkane can be used as an indicator of depositional environment [13,18]. Some biomarker analyses have also been performed by earlier researchers on some source rocks, including the sediment of the Sydney basin [12], the Niger Delta oil [19], and the Ordos Basin oil [20]. But until now there is no organic geochemistry study of the crude oil from Tarakan Basin to determine the depositional environment, maturity and the source of its organic matters. This paper will analyze the aspect of organic geochemistry through the analysis of the aliphatic hydrocarbon biomarker to determine the potency of crude oil from the Tarakan Basin, North Borneo before reactivation.

II. METHOD

Bitumen and water content in oil sample was separated. 2

grams bitumen is separated from the Malten layer in 50 mL *n*-hexane. Malten is fractionated by chromatography column of silica gel GF₂₅₄ based on solvent gradient system. Elution has done with *n*-hexane to obtain an aliphatic hydrocarbon fraction, dichloromethane for obtaining aromatic hydrocarbon fractions, and methanol to obtain a polar fraction [21,22,23]. This Paper only reports aliphatic hydrocarbon fractions. Analysis of the aliphatic hydrocarbon fraction is carried out with the Gas Chromatography-Mass Spectrometer (KG-SM) Agilent GCMS D5975C, a column type HP-5MS (30 m x 250 μ m x 0.2 μ m) with 5% phenyl methyl Silox and helium (He) as the carrier gas. The temperature setting of the column is 70°C (isothermal for 2 minutes) and then increased temperature up to 100°C at a rate of 10°C/min and increased back to 300°C, with a flow rate of 4°C/min and isothermal for 20 minutes. Biomarker of aliphatic hydrocarbon fraction is identified based on specific *m/z* fragmentation, retention time, mass spectrum, and the comparison of previous research results that have been published in the literature.

III. RESULTS AND DISCUSSION

The structure of the aliphatic hydrocarbon biomarker of the Tarakan Basin crude oil is identified based on the specific *m/z* Fragmentogram, retention time and through the comparison of the mass spectrum obtained with the mass spectrum data that has been published by earlier researchers (e.g. Bray and Evan (1961), Eglinton and Hamilton (1967), Seifert and Moldowan (1978), Philp (1985), Lu and Kaplan (1992), Wang et al. (1994), Chattopadhyay and Dutta (2014), Li et al. (2018), Jiang and George (2018)). The identified aliphatic hydrocarbon biomarker consist of a group of *n*-alkanes, isoprenoids, bicyclic sesquiterpenoids, and pentacyclic triterpenoids as seen in Total Ion Chromatogram (TIC) Figure 1 and further spelled out in table 1. The sterane compound is not found in the Tarakan Basin oil samples. The most abundant aliphatic hydrocarbon biomarker is pristane.

A. *n*-Alkane and isoprenoid

The identification of *n*-alkane and isoprenoid compounds against Tarakan oil is conducted based on Fragmentogram *m/z* 57, retention time data and mass spectrum data compared to data that has been reported by past researchers (e.g. Bray and Evan (1961), Eglinton and Hamilton (1967), and Li et al. (2018)). Fragmentogram *n*-alkane and isoprenoid compounds as seen in Figure 2 show homologous *n*-alkane distributed in *n*-C₁₆ – *n*-C₃₀ range with the unimodal distribution type. The short-chain *n*-alkane distribution (under *n*-C₁₆) looks very small, but the isoprenoid Pristane delivers the highest abundance in the oil compound group analyzed in this study. The low intensity of the short-chain *n*-alkane indicates the absence of (very small) algae role in the formation of the Tarakan oil organic compounds. The short-chain *n*-alkane existence (<C₁₅) indicates that organic compounds are derived from phytoplankton, zooplankton, sea bacteria, and algae [6,24]. The identification of the medium chain *n*-alkane group (*n*-C₁₉–*n*-C₂₅) indicates the existence of bacterial activity in the paleoenvironmental of Tarakan oil [25,9,6,26].

The maximum concentration for the medium chain *n*-alkane (*n*-C₁₉ – *n*-C₂₅) is *n*-C₁₉. Previous *n*-C₁₉ existence was also detected with very high concentrations in green algae and anaerobic bacteria [27,28]. The predominance of *n*-C₁₉ was also found in the study of El Nady et al. (2014) on Egyptian

Gulf of Suez Oil and Liu et al. (2019) on the coal samples of the Slovenia basin and indicating that organic matter derived from microbial organisms. The existence of long-chain *n*-alkane (*n*-C₂₆–*n*-C₃₀) with abundance almost equal to medium-chain *n*-alkane (except *n*-C₁₉) indicates the source of organic matter derived from the wax layer of higher terrestrial plants [9]. The same was found in the research of Bakar et al. (2011) with a long chain of *n*-alkane in the *n*-C₂₅ – *n*-C₃₇ range and in the research of Xiaou et al. (2019) with a long chain of *n*-alkane in the *n*-C₂₅ – *n*-C₃₈ range. Both indicate the input of organic matter derived from higher terrestrial plants [9,29].

Odd carbon predominance against even carbon on long-chain *n*-alkane (*n*-C₂₅–*n*-C₃₀) especially on *n*-C₂₅, *n*-C₂₇ and *n*-C₂₉ showed that the organic matter of Tarakan oil were derived from higher terrestrial plants wax layer [9,24,30] that produced through fatty acid decarboxylation [31]. Odd carbon domination of even carbon is also demonstrated by the CPI value of 0.95 as seen in table 3. The value of CPI (Carbon Preference Index) can be used to determine the maturity level of a sample. Dessort (2013) reported CPI value of close to 1 indicated a mature sample, as well as the Onojake et al (2015) study which has CPI value between 0.72 – 1.09. Therefore the value of CPI 0.95 indicates a mature sample of Tarakan oil.

The LHCPI value of 2.03 indicates the relative dominance of the medium chain to the long chain [32]. This is due to the predominance of *n*-C₁₉. This LHCPI value indicates a sufficiently high input from the bacteria [33]. The high *n*-C₁₉ intensity indicates the presence of bacteria that persist in catagenesis stge. The specific bacteria for *n*-C₁₉ are photosynthetic bacteria [34].

The identification based on fragmentogram *m/z* 57 with molecular ion *m/z* 268 indicating the presence of Pristane (Ph) and molecular ion *m/z* 282 indicates the presence of Phytane (Ph) with a ratio of Pr/Ph 3.76 in an analysed oil sample. The value of this Pr/Ph ratio can be used to determine the oil depositional environment. The high-ratio of Pristane/Phytane (Pr/Ph) (>3.0) indicates the input of terrestrial organic matter in oxic conditions whereas the low Pr/Ph (<0.8) indicates an anoxic/hypersaline or carbonated depositional environment [35,36,37]. Therefore, the ratio of Pr/Ph of 3.76 in the sample crude oil from Tarakan Basin indicating that the oil samples were formed in a oxic environment with terrestrial organic input [38].

Besides the ratio of Pr/Ph, organic geochemical aspect of the analyzed Tarakan Basin oil can also be determined based on isoprenoid against *n*-alkanes ratio to know the source of input of organic matter, biodegradation, and maturity level. The Pr/*n*-C₁₇ ratio of the Tarakan Basin oil sample has a value of 34.41 while the Ph/*n*-C₁₈ value is 16.02. When plotted into the chart as shown in Figure 3, it is obtained that the source of organic matter from the sample comes from a higher terrestrial plant with an oxic depositional environment. This data is in accordance with the ratio of Pr/Ph = 3.76 as an indicator of oxic environment of the Tarakan basin which is analyzed and the dominance of odd carbon to even carbon from the homologous series *n*-alkanes. Therefore, this data reinforces the previous analysis of the suspicion that the source of the organic matter of crude oil from Tarakan Basin comes from higher terrestrial plants in oxic depositional environment.

B. Bicyclic sesquiterpenoid

The presence of sesquiterpenoid compounds in Tarakan Basin oil samples was identified based on m/z fragmentogram. Identification of the structure of each compound was based on fragmentation reactions and by comparing the mass spectrum obtained with the published mass spectrum [39,40,41,42,]. The eight peaks in the fragmentogram were identified as C₁₄ bicyclic sesquiterpane, C₁₅ bicyclic sesquiterpane, trans-cadinane, 4β(H)-eudesmane, 8β(H)-drimane, homocadinane, homodrimane, and C₁₆ bicyclic sesquiterpane, where the dominant peaks are C₁₅ and C₁₆- bicyclic sesquiterpane.

The dominance of C₁₅ bicyclic sesquiterpane in the oil samples analyzed indicated that organic matter was derived from higher terrestrial plants resins and prokaryotic bacteria. This assumption is also supported by the presence of C₁₄ sesquiterpane compounds in low intensity [20,43,44,45]. Cadinane is derived from polycadinene precursors found in dammar resin of Angiospermae plants, especially the family of Dipterocarpaceae which undergoes a depolymerization reaction [46], so that the presence of trans-cadinane compounds is considered an indicator of higher Angiospermae plants. In samples there also found 4β(H)-eudesmane where these compounds can be found in fossil plant tissue. The structure of the eudesmane compound is related to the order of terpenoid carbon originating from higher terrestrial plants [6,41,45,47,48]

The presence of 8β(H)-drimane compound together with homodrimane was also identified in the analyzed oil sample. These two compounds identified indicate bacterial input to the formation of organic oil compounds [47]. This is due to the relationship between the structure of the biomarker and the bacteriophanepoliol precursor produced by Archaeobacteria [47], so that the presence of these two compounds can also be associated with bacterial input on the formation of organic matter. The presence of drimane and homodrimane compounds as indicators of bacterial input has also been reported previously in low rank coal samples, Sanggata, East Kalimantan [45], Meghalaya coal samples, India [49] and rock samples in the Gippsland Basin, Australia [48].

The ratio of drimane against homodrimane (as calculated based on Table 1) can be used to determine the depositional environment. The ratio of drimane against homodrimane value of the Tarakan Basin Oil sample is 1.058. According to Weston et al. (1989) and Yangming (2001), this value indicates that the sample is deposited in an oxic environment. The rearrangement mechanism of these two compounds is similar to diasterane [47], which requires acidic clay minerals as catalysts formed in the oxidic depositional environment [50].

The presence of homocadinane in the Tarakan Basin oil, ascribed to the same precursor as cadinane, is in the form of polycadinane as an abundant precursor in high-resin Angiospermae [51]. Therefore, the identification of this compound, potentially used as an indicator of high terrestrial plants, Angiosperms. Similar compounds have also been reported in the oil samples of the Surma Basin, Bangladesh [52] and in the oil samples of the Yinggehai Basin, China [53]. The presence of C₁₆ bicyclic sesquiterpane compounds indicates that oil is formed in the terrestrial environment and the organic matter derived from higher terrestrial plants [41]. This compound is thought to be formed by the degradation of oleanoids from Angiospermae at diagenesis stage [53,54].

C. Pentacyclic Triterpenoid

Identification structure of triterpane pentacyclic compounds in Tarakan oil samples was studied based on m/z 191 fragmentogram and its retention time. Identification of biomarker structures is also done by comparing the mass spectrum obtained with the mass spectrum that has been published by previous researchers (eg Seifert and Maldowan (1978), Philp (1985), Wang et al. (1994), Jiang and George (2018)). The six peaks in the fragmentogram were identified as cis-cis-trans-bicadinane, trans-trans-bicadinane, C₃₀ triterpane, 17α(H),21β(H)-norhopane, 18α(H)-oleanane, and 17α(H),21β(H)-hopane with the most dominant compound is 17α(H),21β(H)-hopane.

The cis-cis-trans-bicadinane and trans-trans-trans-bicadinane compounds identified in the Tarakan Basin oil sample analyzed were cyclization products of cadinene dimeric formed from polycadinene macromolecules through depolymerization and ring closure reactions. Polycadinene has been isolated from both fossils and extracts of the dammar resin of the Angiospermae family of the Dipterocarpaceae family, whose vegetation is abundant in the Tertiary period [46,52,55]. Therefore, the existence of this compound is also potential as an indicator of the Angiospermae family Dipterocarpaceae. The same was reported in the Surma Basin oil sample, Bangladesh [52] and in Northeast Indian sediment samples [39]. In addition, triterpane compounds, generally C₃₀, are also mostly from bacteria. Because the synthesis of triterpenoids in sediments is carried out by microbial, this compound is also used as an important indicator to determine the presence of bacterial input on the formation of organic matter in crude oil [56]. The same was reported by Zhu et al. (2018) in crude oil from the Yinggehai basin and the Qiongdongnan basin, China

The presence of bicyclic compounds identified based on the m/z 191 fragmentogram in the analyzed sample is closely related to one of the bicyclic sesquiterpane compounds namely cadinane. This is because these two compounds come from the same precursor, namely polycadinene [6]. Therefore, the discovery of these two compounds simultaneously confirmed the assumption that the Tarakan Basin oil sample originated from the Angiospermae family Dipterocarpaceae.

The oleanane compound identified in the oil sample analyzed was a triterpenoid pentacyclic compound derived from β-amyrin as a precursor [48,57]. This β-amyrin compound is produced in terrestrial plants Angiosperms whose vegetation is abundant in the Tertiary era [58]. The oleanane compound has 2 isomers namely 18α(H)-oleanane and 18β(H)-oleanane, where the alpha (α) configuration is more stable [59]. Therefore, the presence of 18α(H)-oleanane compound in the oil sample analyzed indicated that the Tarakan Basin oil originated from higher Angiosperm plants and was a mature oil. The existence of this oleanane compound has also been previously reported in rock samples from the Gippsland Basin, Australia [39].

Hopane compounds with configuration of 17α(H), 21β(H)-hopane based on m/z 191 fragmentogram were also identified in the analyzed oil samples. The presence of hopane compounds can be used to determine the maturity of the sample. Biomarker hopane has several isomeric configurations namely 17β(H),21β(H); 17β(H), 21α(H) and 17α(H),21β(H) which are the most stable isomers. The high intensity of the isomers that have high structural stability indicates mature sediment samples. [60,61,62,63].




Figure 1. Total Ion Chromatogram (TIC) aliphatic hydrocarbon fraction of crude oil from Tarakan Basin. Condition: KG-SM temperature program 70°C (2 min), 70°C-100°C (10°C/min), 100°C-300°C (4°C/min) isotherm temperature at 300°C.

Table 1.
Percent Intensity Of Each Peak Of The Aliphatic Hydrocarbon Fraction Of Basin Oil

Group	Retention time	Compound	% intensity
<i>n</i> -alkana	27.09	<i>n</i> -C ₁₆	2.008
	32.77	<i>n</i> -C ₁₇	2.906
	37.25	<i>n</i> -C ₁₈	1.658
	41.49	<i>n</i> -C ₁₉	12.856
	45.53	<i>n</i> -C ₂₀	6.747
	49.63	<i>n</i> -C ₂₁	6.602
	56.47	<i>n</i> -C ₂₂	2.062
	59.07	<i>n</i> -C ₂₃	1.733
	63.11	<i>n</i> -C ₂₄	5.267
	66.99	<i>n</i> -C ₂₅	7.205
	70.76	<i>n</i> -C ₂₆	2.982
	74.42	<i>n</i> -C ₂₇	2.705
	77.97	<i>n</i> -C ₂₈	2.725
	81.24	<i>n</i> -C ₂₉	3.127
	85.4	<i>n</i> -C ₃₀	5.867
isoprenoid	33.21	Pristane	100.000
	37.79	Phytane	26.571
Bicyclic sesquiterpenoid	18.72 , 19.98	C ₁₄ bicyclic sesquiterpane	11.589
	20.98 , 22.88	C ₁₅ bicyclic sesquiterpane	12.533
	22.62	Trans-cadinane	3.651
	23.37	4β(H)-eudesmane	1.720
	23.73 , 24.25	C ₁₆ bicyclic sesquiterpane	13.280
	24.53, 24.67	8β(H)-drimane	7.713
	24.86	Homocadinane	4.464
	28.02	Homodrimane	7.287
Pentacyclic triterpenoid	78.53	cis-cis-trans-bicadinane	1.061
	82.36	trans-trans-trans-bicadinane	1.762
	84.86, 85.08, 85.96	C ₃₀ triterpane	4.470
	86.77	17α(H)-21β(H)-norhopane	3.977
	88.87	18α(H)-oleanane	4.259
	89.22	17α(H)-21β(H)-hopane	6.392




Figure 2. Fragmentogram m/z 57 aliphatic hydrocarbon fraction that indicates *n*-alkanes and isoprenoids

Table 2.
Molecular Parameter For Crude Oil From Tarakan Basin

Parameter	Value
CPI ^a	0.95
LHCPI ^b	2.03
LHCPI ^b	3.76
Pr/ <i>n</i> -C ₁₇	34.41
Ph/ <i>n</i> -C ₁₈	16.02

^a CPI [61] = $\frac{1}{2} \left(\frac{(C_{25}+C_{27}+C_{29}+C_{31}+C_{33})}{(C_{24}+C_{26}+C_{28}+C_{30}+C_{32})} + \frac{(C_{25}+C_{27}+C_{29}+C_{31}+C_{33})}{(C_{26}+C_{28}+C_{30}+C_{32}+C_{34})} \right)$

^b LHCPI = $\frac{C_{17}+C_{18}+C_{19}}{C_{27}+C_{28}+C_{29}}$




Figure 3. Graph plot of Pr/*n*-C₁₇ versus Ph/*n*-C₁₈ of crude oil from Tarakan Basin. Graph plotted based on [68].

Therefore, the identification of compound $17\alpha(\text{H}),21\beta(\text{H})$ -hopane with the highest abundance, but no other hopane isomers are found in the form of $17\beta(\text{H}),21\beta(\text{H})$; $17\beta(\text{H}),21\alpha(\text{H})$ is very strong evidence that the Tarakan Basin oil is mature. The presence of this compound $17\alpha(\text{H}),21\beta(\text{H})$ -hopane has also been reported previously in the sample Brown Shale Central Sumatra Basin[64] and samples of the Central Sichuan Basin, China[65]. The presence of several hopanoid compounds in the analyzed oil samples also shows the presence of bacterial input in the oil formation environment, because hopanoid compounds are produced by bacteriophopanehopanol precursors produced by aerobic bacteria [40,31,62]. The presence of bacterial activity that plays a role in the ancient environment of oil formation in the Tarakan Basin, has also been proven by the presence of other biomarkers in the analyzed samples, such as low-intensity of short chain *n*-alkanes, high intensity of *n*-C₁₉, C₁₄-bicyclic sesquiterpane, drimane and homodrimane. High intensity of *n*-C₁₉ indicates that the bacteria that play a role in Tarakan Basin's paleoenvironmental are photosynthetic bacteria [34].

IV. CONCLUSION

Organic geochemical studies through biomarker analysis of aliphatic hydrocarbon fraction of crude oil from Tarakan Basin, North Borneo provide information of organic matter, depositional environment, and level maturity of the sample. Four groups of identified aliphatic compounds, namely *n*-alkane, isoprenoid, bicyclic sesquiterpenoid and pentacyclic triterpenoid. The presence of long chain *n*-alkanes, trans-cadinane, homocadinane, $4\beta(\text{H})$ -eudesmane, bicadinane, and $18\alpha(\text{H})$ -oleanane indicates organic matter derived from resin dammar Angiospermae family Dipterocarpaceae. It is also reported that there are drimane together with homodrimane and hopane as an indicator of bacterial input. The LHCPI value of 2.03 also indicates a high input of photosynthetic bacteria. The CPI value of 0.95 indicates a mature sample. Pr/Ph ratio of 3.76 and a drimane/homodrimane ratio of 1.058 indicating the oxic depositional environment of the sample. Isomer analysis of $18\alpha(\text{H})$ -oleanane and $17\alpha(\text{H}),21\beta(\text{H})$ -hopane provides information that the crude oil from Tarakan Basin is mature.

ACKNOWLEDGEMENT

The author would like to thank LPPM ITS for providing research funding and to the Molecular Geochemistry Laboratory, Department of Chemistry-Faculty of Science of ITS which has provided laboratory facilities so that this research can be completed.

REFERENCES

- [1] Yudiartono, Anindhita, A. Sugiyono, L. M. A. Wahid, and Adiarso, *Outlook Energi Indonesia 2018: Energi Berkelanjutan untuk Transportasi Darat*. Tangerang: Pusat Pengkajian Industri Proses dan Energi, 2018.
- [2] H. Doust and R. A. Noble, "Petroleum systems of Indonesia," *Marine and Petroleum Geology*, vol. 25, pp. 103–129, 2008.
- [3] Z. Ahmad and L. Samuel, "Stratigraphy and Depositional Cycles in the NE Kalimantan Basin," in *Indonesian Petroleum Association 13th Annual Conference*, 1984, pp. 109 - 120.
- [4] A. Nawawi, A. Suseno, and N. Heriyanto, *Petroleum Geology of Indonesian Basins Volume V Tarakan Basin Northeast Kalimantan*. Jakarta: Pertamina, 1996.
- [5] N. Soeparyono and P. Lennox, "Structural development of hydrocarbon traps in the Cepu oil fields, northeast Java, Indonesia," *Journal of Southeast Asian Earth Sciences*, vol. 4, pp. 281 - 291, 1990.
- [6] K. E. Peters, C. C. Walters, and J. M. dan Moldovan, *The Biomarker Guide: I. Biomarkers and Isotopes in the Environment and Human History*, 2nd ed. USA: Cambridge University Press, 2005.
- [7] K. E. Peters and M. G. Fowler, "Application of petroleum geochemistry to exploration and reservoir management," *Organic Geochemistry*, vol. 33, no. 1, pp. 5 - 36, 2002.
- [8] K. A. Kvenvolden, "Origins of Organic Geochemistry," *Organic Geochemistry*, vol. 39, no. 8, pp. 905 - 909, 2008.
- [9] B. Tissot and D. Welte, *Petroleum Formation and Occurrence*, 2nd ed. Berlin: Springer-Verlag, 1984.
- [10] K. E. Peters, "Guidelines for evaluating petroleum source rock using programmed pyrolysis," *AAPG Bulletin*, vol. 70, pp. 318 - 329, 1986.
- [11] G. Hu, R. Yang, L. Wang, W. Hu, and J. Cao, "Hydrocarbon potential and depositional environment of the Lower Cretaceous black mudstones and shales in the coastal Guangdong Province, China," *Marine and Petroleum Geology*, vol. 99, pp. 92–106, 2019.
- [12] M. Ahmed, H. Volk, S. C. George, M. Faiz, and L. Stalker, "Generation and expulsion of oils from Permian coals of the Sydney Basin, Australia," *Organic Geochemistry*, vol. 40, pp. 810-831, 2009.
- [13] M. A. Al-Atta, G. I. Issa, M. A. Ahmed, and M. M. Afife, "Source Rock Evaluation and Organic Geochemistry of Belayim Marine Oil Field, Gulf of Suez, Egypt," *Egyptian Journal of Petroleum*, vol. 23, pp. 285 - 302, 2014.
- [14] H. Xiao et al., "Oil-oil and oil-source rock correlations in the Muglad Basin, Sudan and South Sudan: New insights from molecular markers analyses," *Marine and Petroleum Geology*, vol. 103, pp. 351 - 365, 2019.
- [15] M. M. El Nady, F. M. Harb, and N. S. Mohamed, "Biomarker characteristics of crude oils from ashrafi and GH oilfields in the Gulf of Suez, Egypt: an Implication to Source Input an Paleoenvironmental Assesments," *Egyptian Journal of Petroleum*, 2014.
- [16] M. C. Onojake, L. C. Osuji, and S. Abrakasa, "Source, depositional environment and maturity levels of some crude oils in southwest Niger Delta, Nigeria," *Chin Journal Geochemistry*, vol. 34, no. 2, pp. 224-232, 2015.
- [17] B. K. G. van Aarssen, T. P. Bastow, R. Alexander, and R. I. Kagi, "Distributions of methylated naphthalenes in crude oils: indicators of maturity, biodegradation and mixing," *Organic Geochemistry*, vol. 30, no. 10, pp. 1213 - 1227, 1999.
- [18] M. H. Hakimi, A. F. Ahmed, and W. H. Abdullah, "Organic Geochemical and Petrographic Characteristics of The Miocene Salif Organic-Rich Shales in The Tihama Basin, Red Sea of Yemen: Implications for Paleoenvironmental Conditions and Oil-Generation Potential," *International Journal of Coal Geology*, pp. 154 - 155, 193 - 204, 2016.
- [19] H. P. Nytoft, G. K. Andersen, T. S. Knudsen, K. Stojanovic, and F. Rise, "Compound 'J' in Late Cretaceous/Tertiary terrigenous oils revisited: Structure elucidation of a rearranged oleanane coeluting on GC with $18\beta(\text{H})$ -oleanane," *Organic Geochemistry*, vol. 77, pp. 89 - 95, 2014.
- [20] L. Ji, C. He, M. Zhang, Y. Wu, and X. Li, "Bicyclic alkanes in source rocks of the Triassic Yanchang Formation in the Ordos Basin and their inconsistency in oil-source correlation," *Marine and Petroleum Geology*, vol. 72, pp. 359 - 373, 2016.
- [21] M. Radke, D. Leythaeuser, and M. Teichmüller, "Relationship between rank and composition of aromatic hydrocarbons for coals of different origins," *Organic Geochemistry*, vol. 6, pp. 423–430, 1984.
- [22] Y. Zetra et al., "Organic geochemistry characteristic of aliphatic hydrocarbon fraction of Sawahlunto coal, Ombilin Basin, West Sumatra," in *The 3rd International Seminar on Chemistry*, Surabaya, 2018, pp. 1-7.
- [23] M. R. Garcia, A. P. Cattani, P. C. Lana, R. C. S. Figueira, and C. C. Martins, "Petroleum biomarkers as tracers of low-level chronic oil contamination of coastal environments: A systematic approach in a subtropical mangrove," *Environmental Pollution*, vol. 249, pp. 1060 - 1070, 2019.

- [24] C. G. Massone, A. de L. R. Wagener, H. M. Abreu, and Á. de Veiga, "Revisiting hydrocarbons source appraisal in sediments exposed to multiple inputs," *Marine Pollution Bulletin*, vol. 73, no. 1, pp. 345 - 354, 2013.
- [25] E. Gelpi, H. Schneider, J. Mann, and J. Oró, "Hydrocarbons of geochemical significance in microscopic algae," *Phytochemistry*, vol. 9, pp. 603 - 612, 1970.
- [26] M. J. Fabiańska and S. Kurkiewicz, "Biomarkers, Aromatic Hydrocarbons and Polar Compounds in the Neogene Lignites and Gangue Sediments of The Konin and Turoszów Brown Coal Basins (Poland)," *International Journal of Coal Geology*, vol. 107, pp. 24 - 44, 2013.
- [27] J. Han, E. D. McCarthy, W. V. Hoeven, M. Calvin, and W. H. Bradley, "Organic geochemical studies, II. A preliminary report on the distribution of aliphatic hydrocarbons in algae, in bacteria, and in a recent lake sediment," in *Proceedings of the National Academy of Sciences of the United States of America*, USA, 1968, pp. 29 - 33.
- [28] G. Li et al., "Microbial production of long-chain n-alkanes: Implication for interpreting sedimentary leaf wax signals," *Organic Geochemistry*, vol. 298, pp. 24 - 31, 2018.
- [29] G. Englington and R. J. Hamilton, "Leaf Epicuticular Wax," *Science*, vol. 156, pp. 1322 - 1335, 1967.
- [30] K. Stojanović and D. Životić, "Comparative study of Serbian Miocene coals — Insights from biomarker composition," *International Journal of Coal Geology*, vol. 107, pp. 3 - 23, 2013.
- [31] S. Killops and V. Killops, *Introduction to organic geochemistry*, 2nd ed. United Kingdom: Blackwell Publishing, 2005.
- [32] A. Aldahik, *Crude Oil Families in the Euphrates Graben Petroleum System*. Berlin: Berlin Institute of Technology, 2011.
- [33] M. G. Commendatore, M. L. Nievas, O. Amin, and J. L. Esteves, "Sources and distribution of aliphatic and polyaromatic hydrocarbons in coastal sediments from the Ushuaia Bay (Tierra del Fuego, Patagonia, Argentina)," *Marine Environment*, vol. 74, pp. 20 - 31, 2012.
- [34] A. El Nemr, A. A. Moneer, S. Ragab, and A. El Sikaily, "Distribution and sources of n-alkanes and polycyclic aromatic hydrocarbons in shellfish of the Egyptian Red Sea coast," *Egyptian Journal of Aquatic Research*, vol. 42, pp. 121 - 131, 2016.
- [35] B. M. Dydik, B. R. T. Simoneit, S.C. Brassell, and G. Eglinton, "Organic geochemical indicators of palaeoenvironmental conditions of sedimentation," *Nature*, vol. 272, pp. 216 - 222, 1978.
- [36] H. L. ten Haven et al., "Application of biological markers in the recognition of palaeo-hypersaline environments," *Fleet, A.J., Kelts, K., Talbot, M.R. (Eds.), Lacustrine Petroleum Source Rocks, Geological Society Special Publication*, vol. 40, pp. 123 - 130, 1988.
- [37] H. K. A. Idris, I. Salihu, M. N. Abdulkadir, and Almustapha, "Application of geochemical parameters for characterization of oil samples using GCMS technique," *International Journal of Physical Sciences*, vol. 3, pp. 152-155, 2008.
- [38] A. Chattopadhyay and S. Dutta, "Higher plant biomarker signatures of Early Eocene sediments of North Eastern India," *Marine and Petroleum Geology*, vol. 57, pp. 51 - 67, 2014.
- [39] L. Jiang and S. C. George, "Biomarker signatures of Upper Cretaceous Latrobe Group hydrocarbon source rocks, Gippsland Basin, Australia: Distribution and palaeoenvironmental significance of aliphatic hydrocarbons," *International Journal of Coal Geology*, vol. 196, pp. 29 - 42, 2018.
- [40] A. Sugiyono, Anindhita, M. A. Boedoyo, and T. dan Adiarso, *Indonesia Energi Outlook 2015 - Pengembangan Energi untuk Mendukung Pembangunan Berkelanjutan*. Jakarta: BPPT, 2015.
- [41] G. Englington and J. Hamilton R, "Leaf Epicuticular Wax," *Science*, vol. 156, pp. 1322 - 1335, 1967.
- [42] K. Stojanovic et al., "Drmno lignite field (Kostolac Basin, Serbia): origin and palaeoenvironmental implications from petrological and organic geochemical studies," *Journal of the Serbian Chemical Society*, vol. 77, no. 8, pp. 1109 - 1127, 2012.
- [43] W. K. Seifert and J. M. Moldowan, "Application of steranes, terpanes and monoaromatics to the maturation, migration and source of crude oils," *Geochimica et Cosmochimica Acta*, vol. 42, pp. 77 - 95, 1978.
- [44] Philp, "Biological markers in fossil fuel production," *Mass Spectrometry Review*, vol. 4, pp. 1 - 54, 1985.
- [45] S. Lu and I. R. Kaplan, "Diterpanes, triterpanes, steranes, and aromatic hydrocarbons in natural bitumens and pyrolysates from aliphatic humic coals," *Gaxhimica er Cosmochimica Acta*, vol. 56, pp. 2761 - 2788, 1992.
- [46] W. Puttman and W. Kalkreuth, "Comparison of hydrocarbon composition in a sequence of humic coals, cannel coals, and oil shales from the Pictou coalfield, Nova Scotia," *Atlantic Geology*, vol. 93-103, p. 25, 1989.
- [47] M. Vliex, H. Hagemann, and W. Puttmann, "Aromatized arborane/fernane hydrocarbons as molecular indicators of floral changes in Upper Carboniferous/Lower Permian strata of the Saar-Nahe Basin, southwestern Germany," *Geochimica et Cosmochimica Acta*, vol. 58, no. 21, pp. 4689 - 4702, 1994.
- [48] Y. Zetra, I. B. Sosrowidjojo, and R. Y. P. Burhan, "Aromatic biomarker from brown coal, Sangatta Coalfield, East Borneo of Middle Miocene to Late Miocene age," *Jurnal Teknologi*, vol. 78, no. 6, pp. 229-238, 2016.
- [49] B. C. van Aarssen, P. Hoogendoorn, and J. W. Leeuw, "A cadinene biopolymer in fossil and extant dammar resins as a source for cadinanes and bicadinanes in crude oils from South East Asia," *Geochimica et Cosmochimica Acta*, vol. 54, pp. 3021 - 3031, 1990.
- [50] R. P. Adams, "Cedarwood oil-Analysis and properties," in *Linskens, H.F., Jackson, J.F. (Eds.), Modern Methods of Plant Analysis, New Series: Oil and Waxes*. Berlin: Springer-Verlag, 1991, pp. 159-173.
- [51] A. Otto and V. Wilde, "Sesqui-, di-, and triterpenoids as chemosystematic markers in extant conifers—a review," *The Botanical Review*, vol. 67, pp. 141-238, 2001.
- [52] Q.-X. Wu, Y.-P. Shi, and Z.-J. Jia, "Eudesmane sesquiterpenoids from the Asteraceae family," *Natural Product Reports*, vol. 23, pp. 699 - 734, 2006.
- [53] R. Alexander, R. Kagi, and R. Noble, "Identification of the Bicyclic Sesquiterpenes Drimane and Eudesmane in Petroleum," *J. Chem. Soc., Chem. Commun*, 1983.
- [54] Zhu Yangming, "Geochemical Characteristics of Different Kinds of Crude Oils in the Tarim Basin, Northwest China," *Chinese Journal of Geochemistry*, vol. 20, no. 1, pp. 73 - 87, 2001.
- [55] B. H. van Aarssen, "The occurrence of polycyclic sesqui-, tri-, and oligoterpenoids derived from a resinous polymeric cadinene in crude oils from southeast Asia," *Geochimica et Cosmochimica*, vol. 56, pp. 1231 - 1246, 1992.
- [56] H. P. Nytoft, O. J. Samuel, G. Kildahl-Andersen, J. E. Johansen, and M. Jones, "Novel C15 sesquiterpanes in Niger Delta oils: structural identification and potential application as a new markers of angiosperms input in light oils," *Organic Geochemistry*, vol. 40, pp. 595 - 603, 2009.
- [57] Y. Zhu, L. Sun, P. Hao, and L. Tuo, "Geochemical composition and origin of Tertiary oils in the Yinggehai and Qiongdongnan Basins, offshore South China Sea," *Marine and Petroleum Geology*, vol. 2018, pp. 139 - 153, 2018.
- [58] D. W. Waples, *Geochemistry in Petroleum Exploration*. Boston: International Human Resources Development Corporation, 1985.
- [59] J. A. Curiale et al., "The central Myanmar (Burma) oil family--composition and implications for source," *Organic Geochemistry*, vol. 22, no. 2, pp. 237 - 255, 1994.
- [60] M. J. Pearson and M. Alam, "Bicadinanes and other terrestrial terpenoids in immature Oligocene sedimentary rocks and a related oil from the Surma Basin, N.E. Bangladesh," *Organic Geochemistry*, vol. 20, no. 5, pp. 539 - 554, 1993.
- [61] C. M. Ekweozor and N. Telnaes, "Oleanane parameter: verification by quantitative study of the biomarker occurrence in sediments of the Nigeria delta," *Organic Geochemistry*, vol. 16, pp. 401 - 413, 1990.
- [62] D. W. Taylor et al., "Biogeochemical evidence for the presence of the angiosperm molecular fossil oleanane in Paleozoic and Mesozoic non-angiospermous fossils," *Paleobiology*, vol. 32, pp. 179 - 190, 2006.
- [63] M. Alberdi and L. Lopez, "Biomarker 18 α (H)-oleanane: a geochemical tool to assess Venezuelan petroleum systems," *Journal of South American Earth Science*, vol. 13, no. 8, pp. 751 - 759, 2000.
- [64] A. Enslinger, A. van Dorsselaer, C. Spyckerelle, P. Albrecht, and G. Ourisson, "Pentacyclic triterpenes of the hopane type as ubiquitous geochemical markers: Origin and Significance," in *Tissot, B./ Bienner. F. (Eds.), Advances in Organic Geochemistry*. Paris: Edition Techcip, 1974, p. 245 = 260.
- [65] R.Y.P. Burhan et al., "Fossil bacterial ecosystem a methane seeps: Origin of organic matter from Be'eri sulfur deposit, Israel.,"

- Geochimica et Cosmochimica Acta*, vol. 66, no. 23, pp. 4085-4101, 2002.
- [66] Y. Han, G. F. John, and T. P. Clement, "Understanding the thermal degradation patterns of hopane biomarker compounds present in crude oil," *Science of the Total Environment*, vol. 667, pp. 792 - 798, 2019.
- [67] W. K. Seifert and J. M. Maldowan, "Paleoreconstruction by biological markers," *Geochimica et Cosmochimica Acta*, vol. 45, pp. 783 - 794, 1980.
- [68] K. J. Safein, T. X. Nguyen, and R. P. Philp, "Organic geochemical and paleoenvironmental characterization of the Brown Shale Formation, Kiliran sub-basin, Central Sumatra Basin, Indonesia," *Organic Geochemistry*, vol. 112, pp. 137 - 157, 2017.
- [69] B. Liu, M. Vrabec, M. Markic, and W. Puttman, "Reconstruction of paleobotanical and paleoenvironmental changes in the Pliocene Velenje Basin, Slovenia, by molecular and stable isotope analysis of lignites," *International Journal of Coal Geology*, vol. 206, pp. 31 - 45, 2019.
- [70] D. Dessort, "Short Course Mineral and Organic Geochemistry," in *TPA Integrated Week*, Surabaya, 2013.
- [71] E. Bray and E. Evans, "Distribution of n-paraffins as a clue to recognition of source beds," *Geochimica et Cosmochimica Acta*, vol. 22, no. 1, pp. 2-15, 1961.
- [72] A. Rudra, S. Dutta, and S. V. Raju, "The Paleogene vegetation and petroleum system in the tropics: A biomarker approach," *Marine and Petroleum Geology*, vol. 86, pp. 38 - 51, 2017.
- [73] R. J. Weston, R. P. Philp, C. M. Sheppard, and A. D. Woolhouse, "Sesquiterpanes, Diterpanes and other Higher Terpanes in Oils from the Taranaki Basin of New Zealand," *Organic Geochemistry*, vol. 14, pp. 405 - 421, 1989.
- [74] B. Cheng, Z. Chen, T. Chen, C. Yang, and T. G. Wang, "Biomarker signatures of the Ediacaran-Early Cambrian origin petroleum from the central Sichuan Basin, South China: Implications for source rock characteristics," *Marine and Petroleum Geology*, vol. 96, pp. 577 - 590, 2018.
- [75] N. A. Bakar, K. S. Tay, N. Y. M. J. B. Omar, Abas M. R., and B. R. T. Simoneit, "The geochemistry of aliphatic and polar organic tracers in sediments from Lake Bera, Malaysia," *Applied Geochemistry*, vol. 26, pp. 1433 - 1445, 2011.